

Considerations in planning vegan diets: Children.

Overview of the review article by VIRGINIA MESSINA, ANN REED MANGELS. *Journal of the American Dietetic Association*. June 2001 v101 i6 p661. COPYRIGHT 2001 American Dietetic Association

Introduction

Although diets of vegan children meet or exceed recommendations for most nutrients, and vegan children have higher intakes of fiber and lower intakes of total fat, saturated fat, and cholesterol than plant and meat eating children, some studies indicate that they may be low in calcium. In addition, biological availability of zinc and iron from plant foods can be low. Protein needs are slightly higher for vegan children but are easily met with a varied diet that provides adequate energy.

Special attention should be given to dietary practices that enhance absorption of zinc and iron from plant foods. Further, good sources of the [omega]-3 fatty acid linolenic acid should be emphasized to enhance synthesis of the long-chain fatty acid docosahexanoic acid.

Dietetics professionals who counsel vegan families should help parents identify good sources of vitamin B-12, riboflavin, zinc, calcium and Vitamin D. (if sun exposure is not adequate) This should not be problematic, due to the growing number and availability of fortified vegan foods that can help children meet all nutrient needs. Therefore, with appropriate food choices, vegan diets can be adequate for children at all ages.

Vegan diets include only plant foods--grains, vegetables, fruits, legumes, nuts, seeds, and vegetable fats. Although the likelihood of nutritional deficiencies increases with more restrictive diets, adequate vegan diets can be planned when parents are committed to appropriate food choices and follow appropriate meal-planning guidelines. According to both The American Dietetic Association [1] and The American Academy of Pediatrics [2], well-planned vegan diets can support normal growth and development in children.

The purpose of this article is to provide an overview of the literature on vegan diets for children, to identify issues that require particular attention, and to offer recommendations for feeding balanced diets to vegan children.

HEALTH ASPECTS OF VEGETARIAN DIETS

Nutrient intakes of vegan children are generally sufficient and sometimes exceed those of plant and meat eating (omnivore) children. Sanders and Manning [3] found that British vegan school-aged children had higher intakes of fiber and all vitamins and minerals except calcium compared to omnivore children.

Older studies conducted with children following more extreme versions of vegan diets should not form the basis for conclusions about adequacy of vegan diets today because most of those studies focused on families on highly restrictive diets not typical of today's more mainstream vegan families. Also, a much larger variety of nutrient-rich and fortified vegan foods are available today. Vegan diets may reduce risk of some chronic diseases of adulthood that have their origins in childhood.

Vegan children have lower intakes of total fat, saturated fat, and cholesterol than omnivore children who typically have intakes of saturated fat and total fat that exceed the levels suggested in the Dietary Guidelines. Vegan children also have higher intakes of fruits and vegetables as evidenced by their higher intakes of vitamin C [3,4]. Finally, vegan diets may expose children to a greater variety of whole plant foods, which may help to establish healthful lifelong eating habits.

ENERGY NEEDS AND GROWTH OF VEGAN CHILDREN

Children with adequate energy intakes will typically grow at a predictable rate. Studies of vegan children have indicated that energy intake is close to the recommended level [4] or to intakes of nonvegetarian controls [9].

One study examined children aged 4 months to 10 years who lived in an intentional community in Tennessee [10]. The majority of children [73%] were vegan from birth; many [76%] regularly used vitamin and mineral supplements. Means for height for age, weight for age, and weight for height were close to the 50th percentile (the average) of the National Center for Health Statistics values for children. Significant differences in height for between vegan children and the reference population were only seen for ages 5 years and younger, with more vegan children tending to be slightly shorter than most children in the reference population (-0.24 cm for age less than 1 year, -2.01 cm for age to 2 years, -2.06 cm for age 3 to 4 years) Overall the average height for age of vegan children were within the normal range and were not indicative of stunting [11].

Although weights for age of the vegan children were slightly below the reference population (-0.09 to -0.37 kg for ages 0 to 8 years), significant differences in weight for age were only seen at age 9 and 10 years when children were 1.11 kg below the National Center for Health Statistics reference population. Sanders [12] reported that British vegan children aged 1 to 18 years had heights, weights, and head and chest circumferences that were within the normal range (between the 3rd and 97th percentile) when compared with standards for non-vegetarian British children.

Boys tended to be slightly shorter and lighter (below the 50th percentile for non-vegetarian British children) whereas girls tended to be slightly lighter. Growth failure, or failure to thrive, commonly defined as weight for age below the 3rd (or 5th) percentile were not seen in these studies [2].

NUTRIENTS OF SPECIAL INTEREST IN DIETS OF VEGAN CHILDREN

Protein

Dietary protein needs of vegan children are generally met when diets contain adequate energy and a variety of plant foods [2,15]. Protein intakes of vegan children have been shown to be similar to those of non-vegetarian children [14] and higher than the standards [4]. Amino acid intakes are also higher than the standards [4].

Plant protein sources include soy products such as soy milk, tofu, tempeh, and textured vegetable protein; other legumes; grains; nuts; and seeds. The Recommended Dietary Allowances (RDA) for protein for children is based on an allowance for reference protein. The RDA for protein is calculated (adjustment) by amino acid composition and digestibility of proteins consumed [5]. The adjustment (new calculation) for the amino acid composition of a vegan diet containing a variety of protein sources is typically 20% more protein for children younger than age 2 years or 10% to 15% for children ages 2 to 6 years. Older children have a lower amino acid requirement per unit protein [5] and generally require only a small adjustment of their protein requirements for amino acid composition [16].

When dietary protein comes mainly from plant foods it is about 85% digestible. A reasonable adjustment to the current RDA for protein to account for this lower digestibility would be about 10% to 15% [17]. To allow for amino acid composition and digestibility of a vegan diet, it appears that protein needs are increased by 30% to 35% for children younger than age 2 years, 20% to 30% for ages 2 to 6 years, and 15% to 20% for children over age 6 years.

Table 1 shows a suggested protein range for vegan children. The range is an additional 2 to 14 g protein daily, which can easily be met in a diet providing adequate energy.

Table 1
Protein recommendations for vegan compared with non-vegan children

Age (yrs)	Weight (kg)	Suggested	Recommended protein	Recommended protein
		protein(g/kg)	intake Vegans(g/day)	intake Non-vegans(g/day)
1-2 (males and females)	11	1.6-1.7	18-19	13
2-3 (males and females)	13	1.4-1.6	18-21	16
4-6 (males and females)	20	1.3-1.4	26-28	24
7-10 (males and females)	28	1.1-1.2	31-34	28
11-14 (males)	45	1.1-1.2	50-54	45
11-14 (females)	46	1.1-1.2	51-55	46
15-18 (males)	66	1.0-1.1	66-73	59
15-18 (females)	55	0.9-1.0	50-55	44

Protein recommendation for children is based on the Recommended Dietary Allowances for age increased to allow for lower amino acid quality and digestibility.

It is generally accepted that combining complementary proteins (additional protein source) at each meal is not necessary to ensure adequate protein intake [1]. In children, however, when beans were added to a corn-based diet, the supplementary effect was somewhat less when the beans were added at intervals of more than 6 hours [16], suggesting that there may be some advantage to eating more than 1 protein source at a meal or within a period of a few hours. From a practical standpoint, because children often eat frequently throughout the day, strict attention to protein combinations at each meal appears to be unnecessary provided a variety of foods are used.

Calcium

Limited data suggest that calcium intakes of vegan children are below current recommendations [3,6,9]. Fulton et al [4] found average calcium intakes among vegan preschoolers to be 39% to 84% of the current Adequate Intake [6]. Claims are commonly made that vegans require less calcium than omnivores, This issue has not been resolved [22-24].

Consumption of soy products, which appears to be fairly typical among vegans, may also favorably affect bone health, Some research shows that soy isoflavones help to prevent breakdown and stimulate growth of bones [25]. However, there is no research on the effects of these factors in vegan children. Because low bone mass and low calcium intake have been related to increased risk of fracture in children and adolescents [26], all vegan children should meet standard recommendations for calcium.

Although oxalates, phytates, and fiber in plant foods decrease calcium availability, research shows that absorption of calcium from many plant foods is excellent. Calcium absorption from low-oxalate vegetables, such as broccoli, kale, and collard greens, ranges from 52% to nearly 59% compared to 32% for milk [27]. Calcium absorption from beans, nuts, and seeds is considerably lower. Limited studies suggest that calcium in fortified foods, including fortified soy milk, is biologically available [28,29].

Table 2 lists calcium content of both fortified and unfortified foods. If children's diets do not contain adequate sources of dietary calcium, supplements may be advisable. Offering supplements between meals may reduce effects of mineral-mineral interactions on absorption.

Table 2**Top sources of calcium, iron, zinc, and vitamin B-12**

Vitamin or mineral	Amount of source
Calcium	mg
Tofu, 1/2 c processed with calcium sulfate	250-390
Orange juice, 8 oz calcium-fortified	300
Soy milk, 8 oz calcium fortified	250-300
Rice milk, 8 oz calcium-fortified	250-300
Dried figs, 5	258
Apple juice, 8 oz calcium-fortified	200
Blackstrap molasses, 1 Tbsp	187
Collard greens, 1/2 c cooked	178
Sesame seeds, 2 Tbsp	176
Tahini, 2 Tbsp	128
Turnip greens, 1/2 c cooked	125
Kale 1/2 c cooked	90
Broccoli, 1/2 c cooked	89
Soybeans, 1/2 c cooked	87
Almond butter, 1/2 Tbsp	86
Textured vegetable protein, 1/2 c prepared	85
Bok choy, 1/2 c cooked	79
Tempeh, 1/2 c	77
Mustard greens, 1/2 c cooked	75
Navy beans, 1/2 c cooked	64
Vegetarian baked beans, 1/2 c	64
Great northern beans, 1/2 c cooked	60
Orange, 1	56
Black beans, 1/2 c cooked	51
Brazil nuts, 2 Tbsp	50
Almonds, 2 Tbsp	50
Iron	mg
Bran flakes, 1 c	11
Tofu, 1/2 c	6.6
Oatmeal, instant, 1 packet	6.3
Cream of wheat, 1/2 c	5.5
Garbanzo beans, 1/2 c cooked	3.4
Blackstrap molasses, 1 Tbsp	3.3
Lentils, 1/2 c cooked	3.2
Pumpkin seeds, 2 Tbsp	2.5
Navy beans, 1/2 c cooked	2.5
Lima beans, 1/2 c cooked	2.2
Swiss chard, 1/2 c cooked	1.9
Black beans, 1/2 c cooked	1.8
Tempeh, 1/2 c	1.8
Spinach, 1/2 c cooked	1.5
Turnip greens, 1/2 c cooked	1.5
Prune juice, 1/2 c	1.5
Apricots, 1/4 c dried	1.4
Beet greens, 1/2 c cooked	1.4
Pasta, enriched, 1/2 c cooked	1.2
Wheat germ, 2 Tbsp	1.2
Raisins, 1/4 c	1.1
Zinc	mg
Bran flakes, 1 c	5
Ready-to-eat breakfast cereal, 3/4 c	1.5 to 4.0
Tempeh, 1/2 c	1.5
Textured vegetable protein, 1/2 c prepared	1.4
Brazil nuts, 2 Tbsp	1.3
Garbanzo beans, 1/2 c cooked	1.3
Tahini, 2 Tbsp	1.3
Lentils, 1/2 c	1.2
Black-eyed peas, 1/2 c cooked	1.1
Peanut butter, 2 Tbsp	1
Split peas, 1/2 c cooked	1
Oatmeal, instant, 1 packet	1
Peas, 1/2 c	1
Corn, 1/2 c	0.9
Vitamin B-12	micro gm
Ready-to-eat breakfast cereal, 3/4 c	1.5 to 6.0
Meat analogs, 1 oz	2.0 to 6.7
Fortified soy milk/rice milk, 8 oz	1.5 to 3
Nutritional yeast, vegetarian support formula, 1 Tbsp	4.0

Vitamin D

Exposing the hands and face to the sun 2 to 3 times per week for 20 to 30 minutes per session appears to provide adequate vitamin D in light-skinned children who live in moderate climates [30]. However, dark-skinned children need longer sun exposure as do children living in cloudy or northern climates.

Vitamin D is poorly supplied by foods; the best sources in most diets are fortified products, of which fortified cow's milk is the most common for US children. Rickets has been seen in some children who do not consume processed or fortified foods, particularly those who live at northern latitudes [31,32]. If sun exposure is not adequate, parents of vegan children should be counseled to choose vitamin D-fortified products. Vitamin D-fortified soy milk and rice milk and breakfast cereals are available. Cereals containing animal-derived vitamin D may not be acceptable to many vegan families, however.

Vitamin B-12

Vitamin B-12 occurs in substantial amounts only in foods derived from animals. Vegan children can obtain adequate vitamin B-12 from fortified foods, including some breakfast cereals, soy beverages, nutritional yeast, and meat analogs [1]. Unfortified nutritional yeasts, sea vegetables, tempeh, and algae are unreliable sources of vitamin B-12 [33,34]. Parents of vegan children have been shown to be aware of the importance of providing their children with reliable sources of vitamin B-12 [3].

Riboflavin

Although riboflavin is sometimes identified as a nutrient of concern in vegan diets, intakes appear to be similar in vegans and omnivores [3,9]. Some brands of soy milk are fortified with riboflavin but most are not, so careful label reading is advised. Other good sources include nutritional yeast, wheat germ, soybeans, mushrooms, leafy green vegetables, avocados, almonds, sea vegetables, fortified cereals, and enriched grains. Whole grains are not especially high in riboflavin. Including small amounts of enriched grains in place of whole grains can help to boost a child's riboflavin intake.

Zinc

The average zinc densities of the diets of vegan and nonvegan children are quite similar [14,35]. Plant sources of zinc include legumes, whole grain pasta, wheat germ, fortified cereals, nuts, and tofu. However, the bioavailability of zinc from plant foods is reduced by phytate (a plant product) in whole grains and legumes.

Although whole grains are high in phytate, these foods are also higher in zinc than refined products so that the total zinc absorbed is greater from whole vs refined grains [36]. Phytate is decreased when whole grain bread is leavened [36] and in the fermentation of soyfoods such as miso and tempeh [37]. Because of these effects on phytate content of foods, zinc absorption may be higher than expected in vegan diets that utilize more leavened bread and fermented soyfoods. Overt zinc deficiency seen in developing 3rd world countries has been observed primarily in children who consume diets high in unleavened grain products and have other predisposing factors for zinc deficiency such as parasitic infection [38,39]. This cannot be considered predictive of zinc status in Western vegan children.

In short-term studies of adults, total zinc absorbed has been shown to be lower on vegetarian diets than on omnivore diets because of lower zinc intake and absorption [40]. Plasma zinc concentrations were also slightly lower although they were within normal ranges. It is possible that some adaptation takes place over the long term. There is some evidence that, in adults, zinc excretion decreases when dietary zinc is low [41,42]. Whether such compensation occurs in children is not known. Assessment of zinc status is complicated by the fact that plasma zinc levels are not sensitive indicators of zinc pools.

Specific attention should be given to identifying zinc-rich foods for vegan children and to dietary factors that enhance zinc absorption. Protein is a promoter of zinc absorption [36,43] based on studies

of individual meals in which total zinc intake was controlled. Emphasizing foods that are good sources of both zinc and protein, such as legumes and nuts, in diets of vegan children can increase both the content and absorption of zinc.

Techniques such as soaking dried beans and discarding the soaking water before cooking and using yeast-leavened bread and fermented soy products like tempeh and miso can also enhance zinc absorption [44]. If a child is following a vegan diet based on **high-phytate** cereals and legumes, zinc supplementation should be considered, especially in early childhood. [45].

Iron

Mean iron intakes of vegan preschoolers (ages 2 to 5 years) have been shown to be above the current RDA in a study [4] where results did not include iron from iron cookware or mineral supplements. In other studies, vegetarian children ages 6 to 12 years, about one-third of whom were vegan, also had iron intakes above the RDA [46] as did vegan children in Britain whose mean iron intakes were more than twice those of nonvegetarian children [3].

Although vegan children typically have iron intakes above recommended levels, non-heme iron has lower absorbability in the intestines.[47]. Only 1 study we know of has reported iron status in vegan children.

Research shows that vegetarian adults typically have lower iron stores than nonvegetarians and that a greater percentage of lacto-ovo-vegetarians (egg & milk vegetarians) have iron stores below the normal range, although most studies indicate that ferritin levels (iron storage levels) are within the normal range. [17,48-52]. Adult vegetarians, including vegans, generally have no higher rates of anemia than those seen in the general adult population [17].

Vitamin C and other organic acids commonly found in vegetables strongly enhance the intestinal absorption of Non-heme iron [53-56]. Vitamin C intakes of vegan children are frequently high [3,4,46] and this, in conjunction with generous levels of dietary iron, may compensate for the lower biological availability of non-heme iron.

Vegan diets contain phytate and other factors found in soy protein and other legumes [57-59], nuts [60], and whole grains that inhibit iron absorption. Subjects do not appear to adapt to increased levels of dietary phytate [48], although some adaptation to diets with low iron bioavailability does occur [61].

Ascorbic acid and other dietary factors can promote iron absorption even in the presence of phytates [53,60,62-64]. Iron bioavailability from soy foods is variable with iron absorption from tempeh and miso being superior to that from soy flour and tofu [65]. The bioavailability of iron from soy milk has not been reported. Leavening of bread reduces the phytate content of wheat and enhances iron absorption [66].

Because iron deficiency anemia is a common nutrition deficiency among all population groups, (vegan and nonvegan) it is important for parents to identify good sources of iron (including iron-fortified foods and iron supplements as needed) in vegan children's diets and to utilize dietary planning principles that maximize intestinal absorption of nonheme iron. Laboratory evaluation of iron status is appropriate if the child's diet is low in iron.

Fiber

The American Academy of Pediatrics recommends an upper limit for fiber in children's diets of 0.5 g/kg body weight per day [2]. The American Health Foundation has proposed the Age Plus 5 guideline for fiber intake in children with age plus 10 g as an upper limit [67]. This closely conforms to the American Academy of Pediatrics recommendations and is a useful guideline for counseling families.

Fiber intakes of vegan children are higher than those of both omnivores and lacto-ovo-vegetarians and may, in some cases exceed recommendations [3]. It is not known if there is any disadvantage to these higher intakes. If overall food intake is low due to early satiety, vegan children may benefit from the consumption of some lower-fiber foods such as refined grains, fruit and vegetable juices, peeled fruits and vegetables, and added fats.

Fat and Essential Fatty Acids

Research suggests that vegan children consume less fat than omnivore and lacto-ovo-vegetarian children [3,68]. Fat intake accounted for 31.5% of energy in 20 lifelong vegan school-aged children [3] and about 30% of energy in 37 lifelong vegan preschoolers [12].

Omnivore children who consume less than 30% of energy from fat appear to grow and develop normally [69,70] although some research suggests inadequate intakes of some nutrients with reduced fat intakes [71]. Higher-fat foods such as nuts, seeds, avocado, soy products, and vegetable oils play an important role in helping vegan children meet energy and nutrient needs.

Vegan diets are generally lacking in the long-chain [varpi]-3 fatty acids docosahexanoic acid (DHA) and eicosapentanoic acid (EPA) (although small amounts may be obtained from sea vegetables) and some studies show low blood levels of these fatty acids in vegan adults [72]. 3-Fatty acid Linolenic acid can be converted to EPA and DHA but efficiency of conversion is reduced in diets high in the 6- Fatty acid linoleic acid [73]. Sanders and Manning [3] found that vegan children had high intakes of linoleic acid compared to linolenic acid with a ratio of 44:1. Although there are no observed risks associated with diets low in long-chain 3 fatty acids, it may be prudent for vegan children to consume diets that enhance conversion of linolenic acid to DHA. Vegan diets for children should include sources of linolenic acid such as ground flax seed, canola oil, walnuts, and soy products and should make use of added vegetable oils that are relatively low in linoleic acid, like olive oil. Because trans-fatty acids may also inhibit synthesis of the long-chain 3-fatty acids, vegetable oils and tub margarine are better choices in vegan diets than stick margarine [74].

COUNSELING PARENTS OF VEGAN CHILDREN

When counseling parents of vegan children, dietary assessment is necessary to determine which foods are excluded from the diet and which foods are acceptable. Some vegans avoid even small amounts of animal foods including additives that may be derived from animals, like the milk protein casein found in most brands of soy cheese. Many of the same issues that influence diet choices among omnivores, such as lack of knowledge of food preparation, lack of time, and economic constraints, are relevant for vegans. New vegans may not be familiar with a wide variety of vegan foods and may not be knowledgeable about where to find these foods and how to prepare them.

It is important to remember that requirements are for essential nutrients, not for specific foods. The Food Guide Pyramid, designed for those persons eating the typical American diet, is not an effective tool for assessment of adequacy of a vegan diet.

AGE-RELATED ISSUES IN COUNSELING PARENTS OF VEGAN CHILDREN

Toddlers and Preschoolers

As toddlers are weaned from breast milk or infant formula, the diet should be evaluated to ensure adequate food sources of calcium, iron, zinc, and vitamin B-12 are included. In some cases supplements may be indicated, Vegan toddlers should be provided with foods of high energy density and be offered 3 meals and 2 to 3 snacks per day. Popular snack ideas for vegan preschoolers include

nut butters on crackers or miniature bagels, fresh fruit, fruit juices, raw vegetables with hummus or tofu dip, dry breakfast cereals, and breads.

From age 18 months to 3 years,

growth rate slows and appetite often decreases. Children may form strong food and eating preferences and are likely to reject unfamiliar foods. Repeated exposure can result in eventual acceptance of foods [75] and parents should offer a variety of foods and model food preferences as toddlers' food preferences have been shown to be markedly affected by the foods offered them [76].

Choking is a concern for toddlers. Foods such as nuts, raw celery and carrots, and popcorn should be avoided or eaten only by older preschoolers with supervision. Food modifications to reduce choking risk include finely grinding nuts, slicing vegetarian hot dogs lengthwise and then crosswise, cutting raw grapes and cherries in half and removing seeds and pits, and quartering cherry tomatoes.

School-Age Children

School introduces a new set of challenges for vegan families. Children may learn for the first time that their diet is considered "alternative" and nutrition education lessons in the classroom may be at variance with the way the child's family eats. Although some children may experience discomfort with eating differently from friends, this is less likely to be a concern as vegetarianism becomes increasingly mainstream. In many schools, school lunch programs offer very little that is acceptable to vegan children and children may need to rely on lunches brought from home.

Adolescents

Despite good appetites, teenagers, who are often on their own for meals, may not always make the most healthful diet decisions. Teenage girls are often concerned with body weight and may restrict energy, making it difficult to meet the high nutrient needs of this stage of life.

Much concern has been raised about the possible relationship between vegetarian diets among teenage girls and eating disorders. A survey conducted in 1986-1987 among Minnesota teenagers showed that vegetarians were much more likely than nonvegetarians to engage in disordered eating behaviors [77]. Martins et al [78] found that vegetarianism was associated with disordered eating only among certain groups. Although data are limited, it is clear that some girls with eating disorders use vegetarian diets as a tool to mask this behavior. Consuming a vegetarian diet does not appear to raise risk for eating disorders, however [79].

Many American teenagers do not consume enough calcium [80] and some data suggest that lactoovo-vegetarian adolescent girls do not meet their calcium needs [17]. If adolescents do not get enough calcium from dietary sources, a supplement may be necessary.

Teenage girls are at high risk in general for iron deficiency anemia [81]. As noted, vegan diets are typically high in iron, but all vegan teenagers should have diets that are adequate in vitamin C to enhance bioavailability of non-heme iron.

A growing phenomenon is the number of vegan teenagers in omnivore households. Dietitians must be prepared to offer support and a great deal of specific guidance because both the adolescents and their parents may be unfamiliar with vegan foods and guidelines for menu planning. It is crucial to elicit support from parents, because teenagers may not plan optimal diets on their own. Parents can show support by learning about both the health benefits of vegan diets and menu-planning guidelines, by purchasing special foods for the vegan adolescent, and by introducing more vegan meals that the whole family enjoys, such as spaghetti with marinara sauce, as well as meals that can be served with and without meat, such as tacos served with meat or textured vegetable protein or stir fries served with

meat or tofu.

Some ideas for foods to stock for vegan teenagers include: dried fruits, trail mix, popcorn, cheeseless pizza, instant hummus mix and pita bread, muffins, juice, nut butters, instant soups, instant bean mixes, bean burritos, veggie burgers, and fortified soy milk in single-serving cartons.

MEAL-PLANNING GUIDELINES FOR VEGAN CHILDREN

Tables 3 and 4 offer guidelines for feeding vegan children.

Dietitians may need to give parents additional advice for making appropriate and practical choices from within the food groups.

Table 3

Food guide for vegan toddlers

Food	Servings per day	Servings Size
Grains	6+	½ to 1 slice bread; ¼ to ½ c cooked cereal, grain or pasta; ½ - 1c ready-to-eat cereal
Vegetables	2+	¼ to ½ c cooked; ½ to 1 c raw
Fruit	3+	¼ to ½ c canned; ½ c juice; ½ medium piece of fruit
Legumes/nuts	2+ (include at least 1 Serving per day of nuts/seeds or 1 full-fat soy product)	¼ - ½ c cooked beans; tofu, tempeh, or textured vegetable protein; ½ --3 oz meat analog; ½ Tbsp nuts, seeds, or nut or seed butter
Fortified soymilk or breast milk	3	1 c
Fats	3-4	1 tsp margarine or oil

Table 4

Food guide for vegan preschoolers and school-aged children

Food	Servings for for ages 5-6 years	Servings for ages 7-12 years	Serving Size
Grains	6+	7+	Grains 1 slice bread; 1/2 c cooked cereal, grain, or pasta; 3/4-1 c ready- toast cereal
Vegetables	2+	3+	1/2 c cooked; 1 c raw
Fruits	2+	3+	1/2 c canned fruit; 3/4 c juice; 1 medium fruit
Legumes	1-2	2-3	1/2 c cooked beans, tofu, tempeh, or textured vegetable protein; 3 oz meat analog
Nuts	1	1	2 Tbsp nuts, seeds, or nut or seed butter
Fortified soymilk	3	3	1 c
Fats	4	5	1 tsp margarine or oil

Vegetables and Fruits

All children should consume a wide variety of vegetables and fruits. Green leafy vegetables, often rich in calcium and riboflavin, are especially valuable in vegan diets.

Grains

Although the emphasis should be on whole grains, refined grains can help young vegan children meet energy and nutrient needs. Vegan children may come closer to the American Academy of Pediatrics recommendations for fiber intake by consuming half of their servings of grains as refined products.

Legumes

Dried beans, peas, lentils, and soy products play an important role in vegan diets as key sources of energy, protein, iron, zinc, and calcium. Many soy products are especially rich in protein and sometimes calcium and can be versatile and useful as meat substitutes. To encourage children to consume legumes, make legumes into spreads such as hummus and serve on crackers or slices of

apples; serve roasted soynuts as snacks or in salads; or add chopped legumes to loaves or veggie burgers.

Nuts and Seeds

Nuts and seeds are important in the diets of vegan children as they are rich in protein, essential fats, energy, and trace minerals. Spread nut and seed butters on bread, crackers, and slices of raw fruits and vegetables.

Fortified Soy milk

Although soy milk is not essential in the diets of vegan children, the fortified brands are useful in helping children meet needs for calcium and vitamin D. For some families, the cost of soy milk per day may be prohibitive. The use of powdered soy milk (sometimes available in bulk in natural foods stores) may be helpful. When vegan children consume fewer than 3 servings per day, dietitians will need to help parents identify other foods that can help children meet needs for protein, calcium, vitamin D (where necessary) and vitamin B-12. If lower-protein beverages such as fortified rice milk or fruit juices are used in place of soy milk, other protein sources such as legumes, nuts, or grains, should be increased to compensate for the lower protein intake.

Fats

Small amounts of added fats help vegan children meet needs for energy and essential fatty acids. Vegetable oils that are rich in linolenic acid and/or low in linoleic acid are the best choices for regular use. Children can also have moderate amounts of soft margarine to help meet energy needs.

APPLICATIONS/CONCLUSIONS

Vegan diets can meet the nutrition needs of children if appropriately planned by a knowledgeable adult. Limited research indicates that vegan diets can support growth that falls within normal ranges. The wide availability of convenient vegan foods, many of which are fortified, make it increasingly easy to plan healthful vegan diets for children. All vegan children should regularly consume foods fortified with vitamin B-12 or use a supplement. A vitamin D supplement is necessary if sun exposure is not adequate and parents of vegan children need to give extra attention to including adequate calcium, zinc, and iron in the diet and to incorporating dietary practices that enhance absorption of zinc and iron from plant foods. In addition to staying aware of the nutrition issues that are relevant to planning vegan diets, dietetics professionals may need to be familiar with availability of vegan foods in their community and with methods of preparation.

V. Messina is a nutrition consultant with Nutrition Matters, Inc, Port Townsend, Wash. She is also adjunct assistant professor at Loma Linda University, Loma Linda, Calif. A. R. Mangels is a nutrition advisor for the Vegetarian Resource Group, Baltimore, Md.

Additional Nutrient Table from AAP

Table 2. Nutrients and Food Sources

Pediatrics in Review Vol.30 No.1 January 2009 e3

Protein

Tofu, tempeh, legumes, grains, eggs, dairy

Omega-3 Fatty Acids

Flax seed, dark greens, tofu, fish oil, nuts

Iron

Legumes, nuts, dried fruit, spinach, fortified grains

Calcium

Kale, broccoli, fortified orange juice, fortified soy, figs, dairy

Zinc

Whole grains, legumes, nuts, wheat germ, whole grain pasta

Folate

Legumes, dark green leafy vegetables, fortified cereals/breads

Vitamin B12

Fortified eggs, fortified dairy, cereals, breads, some fortified soy

Summary Pediatrics in Review Vol.30 No.1 January 2009 e3

Vegetarian diets can provide appropriate and adequate nutrition for all stages of life.

The key to success is accurate information and understanding of the subject matter.

Because vegetarianism is a complex issue influenced by many different factors, it can be difficult to stay well informed.

Numerous studies have shown several important health benefits for adult vegetarians.

The job of pediatricians is to ensure that their pediatric patients reap the same health benefits as adults.

- Based on strong research evidence, breastfeeding vegetarians should ensure that they are consuming adequate amounts of vitamin B12, folate, and omega 3 fatty acids. (4)(5)(6)
- Based on strong research evidence, vegetarian children raised on a varied diet have normal growth and development measures. (7)(8)(9)(10)(11)(12)(16)
- Based on strong research evidence, adolescent vegetarians choose vegetarianism for very different reasons than their adult counterparts, and their decision often is impulsive and without much forethought. (13)(14)
- Based on strong research evidence, a food guide pyramid that is specific to vegetarians and takes into account the many subtleties of vegetarianism is important when counseling vegetarian patients. (15)
- Based on strong research evidence, vegetarianism in adolescents can be a means of concealing an underlying eating disorder. (13)

References

- (1.) Position of The American Dietetic Association: Vegetarian diets. J Am Diet Assoc. 1997;97:1317-1321.
- (2.) American Academy of Pediatrics, Committee on Nutrition. Pediatric Nutrition Handbook. 4th ed. Elk Grove Village, III: American Academy of Pediatrics; 1998.
- (3.) Sanders TAB, Manning J. The growth and development of vegan children. J Hum Nutr Diet. 1992;5:11-21.
- (4.) Fulton JR, Hutton CL, Stitt KR. Preschool vegetarian children. J Am Diet Assoc. 1980;76:360-365.
- (5.) Food and Nutrition Board. Recommended Dietary Allowances. 10th ad. Washington, DC: National Academy Press; 1989.

(6.) Institute of Medicine, Food and Nutrition Board. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. Washington, DC: National Academy Press; 1997. HWRC Document <http://galenet.galegroup.com.lp.hscl.ufl.edu/servlet/HWRC/hits?docNum...>

10 of 18 1/12/2010 11:16 PM

(7.) Institute of Medicine, Food and Nutrition Board. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B-6, Folate, Vitamin B-12, Pantothenic Acid, Biotin, and Choline. Washington, DC: National Academy Press; 1998

(8.) Institute of Medicine, Food and Nutrition Board. Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids. Washington, DC: National Academy Press; 2000.

(9.) Sanders TAB, Purves R. An anthropometric and dietary assessment of the nutritional status of vegan preschool children. *J Hum Nutr.* 1981;35:349-357.

(10.) O'Connell JM, Dibley MJ, Sierra J, Wallace B, Marks JS, Yip R. Growth of vegetarian children: the Farm Study. *Pediatrics.* 1989;84:475-481.

(11.) Torun B, Chew F. Protein-energy malnutrition. In: Shils ME, Olson JA, Shike M, Ross AC, eds. *Modern Nutrition in Health and Disease.* 9th ed. Baltimore, Md: Williams & Wilkins; 1999:963-988.

(12.) Sanders TAB. Growth and development of British vegan children. *Am J Clin Nutr.* 1988; 48:822-825.

(13.) Jacobs C, Dwyer JT. Vegetarian children: appropriate and inappropriate diets. *Am J Clin Nutr.* 1988;48(suppl):811-818.

(14.) Sanders TAB. Vegetarian diets and children. *Pediatr Clin N Am.* 1995;42:955-965.

(15.) Millward DJ. The nutritional value of plant-based diets in relation to human amino acid and protein requirements. *Proc Nutr Soc.* 1999;58:249-260.

(16.) Young VR, Pellett FL. Plant proteins in relation to human protein and amino acid nutrition. *Am J Clin Nutr.* 1994;59(suppl):1203S-1212S.

(17.) Messina MJ, Messina VL. *The Dietitian's Guide to Vegetarian Diets: Issues and Applications.* Gaithersburg, Md: Aspen; 1996.

HWRC Document <http://galenet.galegroup.com.lp.hscl.ufl.edu/servlet/HWRC/hits?docNum...>

11 of 18 1/12/2010 11:16 PM

(18.) Frassetto LA, Todd KM, Morris RC Jr, Sebastian A. Estimation of net endogenous noncarbonic acid production in humans from diet potassium and protein contents. *Am J Clin Nutr.* 1998;68:576-583.

(19.) Dwyer JT, Foulkes E, Evans M, Ausman L. Acid/alkaline ash diets: time for assessment and change. *J Am Diet Assoc.* 1985;85:841-845.

(20.) Linkawiler HM, Zemel MB, Hegsted M, Schuette S. Protein-induced hypercalciuria. *Fed Proc.* 1981;40:2429-2433.

(21.) Schuette SA, Linkawiler HM. Effects of Ca and P metabolism in humans by adding meat, meat plus milk, or purified proteins plus Ca and P to a low protein diet. *J Nutr.* 1982;112:338-349.

(22.) Massey LK. Does dietary protein adversely affect bone? Symposium overview. *J Nutr.* 1998;128:1048-1050.

(23.) Barzel US, Massey LK. Excess dietary protein can adversely affect bone. *J Nutr.* 1998;128:1051-1053.

(24.) Heaney RP. Excess dietary protein may not adversely affect bone. *J Nutr.* 1998;128:1054-1057.

(25.) Ishida H, Uesugi T, Kuniaki H, Toda T, Nukay H, Yokotsuka K, Tsuji K. Preventive effects of plant isoflavones, daidzin and genistin, on bone loss in ovariectomized rats fed a calcium-deficient diet. *Biol Pharm Bull.* 1998;21:62-66.

(26.) Goulding AR, Williams SM, Gold EJ, Taylor RW, Lewis-Barned NJ. Bone mineral density in girls with forearm fractures. *J Bone Miner Res.* 1998;13:143-148.

(27.) Weaver CM, Plawecki KL. Dietary calcium: adequacy of a vegetarian diet. *Am J Clin Nutr.* 1994;59(suppl):1238S-1241S.

(28.) Andon MB, Peacock M, Kanerva RL, DeCastro JA. Calcium absorption from apple and orange juice fortified with calcium citrate malate (CCM). *J Am Coll Nutr.* 1996;15:313-316.

(29.) Heaney RP, Dowell MS, Rafferty K, Bierman J. Bioavailability of the calcium in fortified soy

- imitation milk, with some observations on method. *Am J Clin Nutr.* 2000;71:1166-1169.
- (30.) Specker BL, Valanis B, Hertzberg V, Edwards N, Tsang RC. Sunshine exposure and serum 25-hydroxyvitamin D concentrations in exclusively breast-fed infants. *J Pediatr.* 1985;107:372-376.
- (31.) Dagnaliev PC, Vergote FJ, van Staveren WA, van den Berg H, Dingjan PG, Hautvast JG. High prevalence of rickets in infants on macrobiotic diets. *Am J Clin Nutr.* 1990;51:202-208.
- (32.) Dwyer JT, Dietz WH Jr, Hass G, Suskind R. Risk of nutritional rickets among vegetarian children. *Am J Dis Child.* 1979;133:134-40.
- (33.) Herbert V. Vitamin B-12: plant sources, requirements, and assay. *Am J Clin Nutr.* 1988;48:852-858.
- (34.) Dagnaliev PC, van Staveren WA, van den Berg H. Vitamin B-12 from algae appears not to be bioavailable. *Am J Clin Nutr.* 1991;53:695-697.
- HWRC Document <http://galenet.galegroup.com.lp.hscl.ufl.edu/servlet/HWRC/hits?docNum...>
12 of 18 1/12/2010 11:16 PM
- (35.) Gibson RS. Content and bioavailability of trace elements in vegetarian diets. *Am J Clin Nutr.* 1994;59(suppl):1223S-1225S.
- (36.) Sandstrom B, Arvidsson B, Cederblad A, Bjorn-Rasmussen E. Zinc absorption from composite meals I. The significance of wheat extraction rate, zinc, calcium, and protein content in meals based on bread. *Am J Clin Nutr.* 1980;33:739-745.
- (37.) Hirabayashi M, Matsui T, Yano H. Fermentation of soybean flour with *Aspergillus ussuriensis* improves availabilities of zinc and iron in rats. *J Nutr Sci Vitaminol.* 1998;44:877-886.
- (38.) Prasad AS, Schulert AR, Sandstead HH. Zinc and iron deficiencies in male subjects with dwarfism but without acrodermatitis, schistosomiasis, or severe anemia. *Am J Clin Nutr.* 1963;12:437-444.
- (39.) Ronaghy HA, Reinhold JG, Mahloudji M, Ghavami F, Fox MRS, Halstead JA. Zinc supplementation of malnourished schoolboys in Iran: increased growth and other effects. *Am J Clin Nutr.* 1974;27:112-121.
- (40.) Hunt JR, Matthys LA, Johnson LK. Zinc absorption, mineral balance, and blood lipids in women consuming controlled lacto-ovo-vegetarian and omnivorous diets for 8 wk. *Am J Clin Nutr.* 1998;67:421-30.
- (41.) Sian L, Mingyan X, Miller LV, Tong L, Krebs NP, Hambidge KM. Zinc absorption and intestinal losses of endogenous zinc in young Chinese women with marginal zinc intakes. *Am J Clin Nutr.* 1996;63:348-353.
- (42.) Campbell-Brown M, Ward RJ, Haines AP, North WRS, Abraham R, McFadyen IR. Zinc and copper in Asian pregnancies-is there evidence for a nutritional deficiency? *Br J Obstet Gynaecol.* 1985;92:875-885.
- (43.) Hunt JR, Lykken GI, Mullen LK. Moderate and high amounts of protein from casein enhance human absorption of zinc from whole wheat or white rolls. *Nutr Res.* 1991;11:413-418.
- (44.) Gibson RS, Yeudall F, Drost N, Mtitimuni B, Cullinan T. Dietary interventions to prevent zinc deficiency. *Am J Clin Nutr.* 1998; 68(suppl):484S-487S.
- (45.) Allen LH. Zinc and micronutrient supplements for children. *Am J Clin Nutr.* 1998; 68(suppl):495S-498S.
- (46.) Kim Y-C. The effect of vegetarian diet on the iron and zinc status of school-age children. [master's thesis]. Amherst: University of Massachusetts; 1986.
- (47.) Hunt JR, Roughead ZK. Nonheme-iron absorption, fecal ferritin excretion, and blood indexes of iron status in women consuming controlled lacto-ovo-vegetarian diets for 8 wk. *Am J Clin Nutr.* 1999;69:944-952.
- (48.) Brune M, Rossander L, Hallberg L. Iron absorption: no intestinal adaptation to a high-phytate diet. *Am J Clin Nutr.* 1989;49:542-545.
- (49.) Ball MJ, Bartlett MA. Dietary intake and iron status of Australian vegetarian women. *Am J Clin Nutr.* 1999;70:353-358.
- HWRC Document <http://galenet.galegroup.com.lp.hscl.ufl.edu/servlet/HWRC/hits?docNum...>
13 of 18 1/12/2010 11:16 PM
- (50.) Alexander D, Ball MJ, Mann J. Nutrient intake and haematological status of vegetarians and

- age-sex matched omnivores. *Eur J Clin Nutr.* 1994;48:538- 546.
- (51.) Halman AD, Darnton-Hill I. Vitamin and iron status in new vegetarians. *Am J Clin Nutr.* 1987;45:785-789.
- (52.) Donovan UM, Gibson RS. Iron and zinc status of young women aged 14- 19 years consuming vegetarian and omnivorous diets. *J Am Coll Nutr.* 1995;14:463-472.
- (53.) Hallberg L, Brune M, Rossander, L. Effect of ascorbic acid on iron absorption from different types of meals. Studies with ascorbate-rich foods and synthetic ascorbic acid given in different amounts with different meals. *Hum Nutr Appl Nutr.* 1986;40:97-113.
- (54.) Cook JD, Mosen ER. Vitamin C, the common cold, and iron absorption. *Am J Clin Nutr.* 1977;30:235-241.
- (55.) Mosen ER, Hallberg L, Layrisse M, Hegated DM, Cook JD, Marts W, Finch CA. Estimation of available dietary iron. *Am J Gun Nutr.* 1978;31:134 141.
- (56.) Gillooly M, Bothwell TH, Torrance JD, MacPhail AP, Derman DP, Bezwoda WR, Mills W, Chariton RW. The affects of organic acids, phytatas and polyphenols on the absorption of iron from vegetables. *Br J Nutr.* 1983;49:331-342.
- (57.) Hallberg L, Rossander L. Effect of soy protein on nonheme iron absorption in man. *Am J Clin Nutr.* 1982;36:514-520.
- (58.) Hurrall RF, Juillerat M-A, Reddy MB, Lynch SR, Dassanko SA, Cook JD. Soy protein, phytata, and iron absorption in humans. *Am J Clin Nutr.* 1992;56:573-578.
- (59.) Lynch SR, Beard JL, Dassenko SA, Cook JD. Iron absorption from legumes in humans. *Am J Clin Nutr.* 1984;40:42-47.
- (60.) Macfarlane BJ, Bexwoda WR, Bothwell TH, Baynes RD, Bothwell JE, MacPhail AF, Lamparelli RD, Mayet F. Inhibitory effect of nuts on iron absorption. *Am J Clin Nutr.* 1988;47:270-274.
- (61.) Hunt JR, Roughead ZK. Adaptation of iron absorption in men consuming diets with high or low iron bioavailability. *Am J Clin Nutr.* 2000;71:94-102.
- (62.) Hallberg L, Brune M, Rossander L. Iron absorption in man: ascorbic acid and dose-dependent inhibition by phytate. *Am J Clin Nutr.* 1989;49:140-144.
- (63.) Siegenberg D, Baynes RD, Bothwell TH, Macfarlane BJ, Lamparelli RD, Car NG, MacPhail P, Schmidt U, Tal A, Mayet F. Ascorbic acid prevents the dose-dependent inhibitory effects of polyphenols and phytates on nonheme iron absorption. *Am J Clin Nutr.* 1991;53:537-541.
- (64.) Hallberg L, Huithen L. Prediction of dietary iron absorption: an algorithm for calculating absorption and bioavailability of dietary iron. *Am J Clin Nutr.* 2000;71:1147-1160.
- (65.) Macfarlane BJ, van der Riet WB, Bothwell TH, Baynes RD, Siegenberg D, Schmidt U, Tol A, Taylor JRN, Mayet F. Effect of traditional oriental soy products on iron absorption. *Am J Clin Nutr.* 1990;51:873-880.
- HWRC Document <http://galenet.galegroup.com.lp.hscl.ufl.edu/servlet/HWRC/hits?docNum...>
14 of 18 1/12/2010 11:16 PM
- (66.) El-Guindi M, Lynch SR, Cook JD. Iron absorption from fortified flat breads. *Br J Nutr.* 1988;59:205-213
- (67.) Williams CL, Bollella M. Is a high fiber diet safe for children? *Pediatrics.* 1995;96(suppl)1014S-1019S.
- (66.) Lenfant C, Ernst N. Daily dietary fat and total food energy intakes--Third National Health and Nutrition Examination Survey, phase 1, 1988-1991. *MMWR Morbid Mortal Wkly Rep.* 1994;43:116-125.
- (69.) Shea S, Basch C, Stein A, Contento I, Irigoyen M, Zybert P. Is there a relationship between dietary fat and stature or growth in children three to five years of age? *Pediatrics* 1993;92:579-586.
- (70.) Boulton TJC, Magarey AM Effects of differences in dietary fat on growth, energy and nutrient intake from infancy to eight years of age *Acta Paediatr.* 1995;84:146-150.
- (71.) Nicklas TA, Webber LS, Koschak M, Berenson GS. Nutrient adequacy of low fat intakes for children: the Bogalusa Heart Study. *Pediatrics.* 1992;89:221-228.
- (72.) Sanders TAB, Ellis FR, Dickerson, JWT. Studies of vegans: the fatty acid composition of plasma choline phosphoglycerides, erythrocytes, adipose tissue, and breast-milk and some indicators of susceptibility to ischaemic heart disease in vegans and omnivore controls. *Am J Clin Nutr.*

Bok choy, 1/2 c cooked	79
Tempeh, 1/2 c	77
Mustard greens, 1/2 c cooked	75
Navy beans, 1/2 c cooked	64
Vegetarian baked beans, 1/2 c	64
Great northern beans, 1/2 c cooked	60
Orange, 1	56
Black beans, 1/2 c cooked	51
Brazil nuts, 2 Tbsp	50
Almonds, 2 Tbsp	50
Iron	mg
Bran flakes, 1 c	11
Tofu, 1/2 c	6.6
Oatmeal, instant, 1 packet	6.3
Cream of wheat, 1/2 c	5.5
Garbanzo beans, 1/2 c cooked	3.4
Blackstrap molasses, 1 Tbsp	3.3
Lentils, 1/2 c cooked	3.2
Pumpkin seeds, 2 Tbsp	2.5
Navy beans, 1/2 c cooked	2.5
Lima beans, 1/2 c cooked	2.2
Swiss chard, 1/2 c cooked	1.9
Black beans, 1/2 c cooked	1.8
Tempeh, 1/2 c	1.8
Spinach, 1/2 c cooked	1.5
Turnip greens, 1/2 c cooked	1.5
Prune juice, 1/2 c	1.5
Apricots, 1/4 c dried	1.4
Beet greens, 1/2 c cooked	1.4
Pasta, enriched, 1/2 c cooked	1.2
Wheat germ, 2 Tbsp	1.2
Raisins, 1/4 c	1.1
Zinc	mg
Bran flakes, 1 c	5
Ready-to-eat breakfast cereal, 3/4 c	1.5 to 4.0
Tempeh, 1/2 c	1.5
Textured vegetable protein, 1/2 c prepared	1.4
Brazil nuts, 2 Tbsp	1.3
Garbanzo beans, 1/2 c cooked	1.3
Tahini, 2 Tbsp	1.3
Lentils, 1/2 c	1.2
Black-eyed peas, 1/2 c cooked	1.1
Peanut butter, 2 Tbsp	1
Split peas, 1/2 c cooked	1
Oatmeal, instant, 1 packet	1
Peas, 1/2 c	1
Corn, 1/2 c	0.9
Vitamin B-12	micro gm
Ready-to-eat breakfast cereal, 3/4 c	1.5 to 6.0
Meat analogs, 1 oz	2.0 to 6.7
Fortified soymilk/rice milk, 8 oz	1.5 to 3
Nutritional yeast, vegetarian support formula, 1 Tbsp	4.0

Table 3
Food guide for vegan toddlers

Food	Servings per day	Servings Size
Grains	6+	1/2 to 1 slice bread; 1/4 to 1/2 c cooked cereal, grain or pasta; 1/2 - 1c ready-to-eat cereal
Vegetables	2+	1/4 to 1/2 c cooked; 1 c raw

Fruit	3+	¼ to ½ c canned; ½ c juice; ½ medium piece of fruit
Legumes/nuts	2+ (include at least 1 Serving per day of nuts/seeds or 1 full-fat soy product)	¼ - ½ c cooked beans; tofu, tempeh, or textured vegetable protein; ½ --3 oz meat analog; ½ Tbsp nuts, seeds, or nut or seed butter
Fortified soymilk or breast milk	3	1 c
Fats	3-4	1 tsp margarine or oil

Table 4
Food guide for vegan preschoolers and school-aged children

Food	Servings for for ages 5-6 years	Servings for ages 7-12 years	Serving Size
Grains	6+	7+	Grains 1 slice bread; 1/2 c cooked cereal, grain, or pasta; 3/4-1 c ready- toast cereal
Vegetables	2+	3+	1/2 c cooked; 1 c raw
Fruits	2+	3+	1/2 c canned fruit; 3/4 c juice; 1 medium fruit
Legumes	1-2	2-3	1/2 c cooked beans, tofu, tempeh, or textured vegetable protein; 3 oz meat analog
Nuts	1	1	2 Tbsp nuts, seeds, or nut or seed butter
Fortified soymilk	3	3	1 c
Fats	4	5	1 tsp margarine or oil